

POSITIVE DEFINITE MAPS, REPRESENTATIONS AND FRAMES

DORIN ERVIN DUTKAY

ABSTRACT. We present a unitary approach to the construction of representations and intertwining operators. We apply it to the C^* -algebras, groups, Gabor type unitary systems and wavelets. We give an application of our method to the theory of frames, and we prove a general dilation theorem which is in turn applied to specific cases, and we obtain in this way a dilation theorem for wavelets.

CONTENTS

1. Introduction	1
2. Positive definite maps and representations	2
3. Intertwining operators	6
4. Frames and dilations	11
5. A dilation theorem for wavelets	14
References	20

1. INTRODUCTION

Engineering problems in time-frequency analysis of coherent vector expansions, Gabor bases, wavelets based on scaling and integral translations, and multiresolution algorithms in signal processing are generally not thought to be related to operator algebras. In this paper, we show nonetheless that a fundamental idea of Kolmogorov adds clarity to known constructions in operator algebra theory, and moreover is the key to an extension of recent results in the more applied areas that we enumerated above. Of our original results (see section 5 below) we highlight a new algorithm for the construction of certain orthonormal frames of wavelet type. Our paper proposes a general method of construction of representations of various algebraic structures as operators on Hilbert spaces. Our goal is to show how some well known constructions of representations fit into the same framework and are consequences of a general result. Among the structures considered, we mention C^* -algebras, groups, Gabor type unitary systems and wavelet representations.

In operator theory, the GNS construction producing representations of C^* -algebras is a fundamental tool (see [BraRo]). In harmonic analysis unitary representations of groups can be constructed when a function of positive type is present (see [Fol]). Representations are ubiquitous also in the theory of wavelets and frames (see [HL], [Jor98]). We will see how these various results have in fact a common ground - a classical theorem of Kolmogorov (theorem 2.2), also known in the literature as the Kolmogorov decomposition of positive definite kernels. We follow here the ideas introduced in [EvLe]. It is shown there that the Kolmogorov theorem gives a unified treatment of several important dilation theorems such as the GNS-Stinespring construction for C^* -algebras, the Naimark-Sz. Nagy unitary dilation of positive definite functions on groups, the construction of Fock spaces and the algebras of canonical commutation and anticommutation relations. Kolmogorov's result was used also by Sz. Nagy and C. Foias in dilation theory, for the commutant lifting theorem ([SzF68], [SzF70]) which in turn was a key idea used by D. Sarason to obtain a solution to the Nevanlinna-Pick interpolation problem ([Sar67]). For a more complete account of the history and applications of Kolmogorov's result, we refer to [C96].

We will indicate how this technique can be used also for construction of wavelet representations and Gabor type unitary systems.

More general constructions for Hermitian kernels are also possible and they are based on Krein spaces (see [C97]).

In section 2 we review the general result of Kolmogorov and we show how it can be used for the GNS construction and for positive definite maps on groups. Then we apply it to Gabor type unitary systems and we obtain unitary representations and for wavelets we get the cyclic representations introduced in [Jor98].

Section 3 concerns operators compatible with the representations defined in section 2, called intertwining operators. Again, the starting point is a general theorem (theorem 3.2). We consider some particular cases and study how the intertwining operators will be compatible with the additional structure that appears.

In section 4 we analyze some connections between representations and frames. We recall that a set $\{x_n \mid n \in \mathbb{N}\}$ of vectors in a Hilbert space H is called a frame for the Hilbert space H if there are some positive constants A and B such that

$$A\|f\|^2 \leq \sum_{n \in \mathbb{N}} |\langle f \mid x_n \rangle|^2 \leq B\|f\|^2, \quad (f \in H).$$

When $A = B = 1$ then we call it normalized tight frame.

It is known that any normalized tight frame is the projection of an orthonormal basis of a bigger Hilbert space (see [HL]). We will prove that the normalized tight frames can be dilated to orthonormal bases in a way that is compatible with the representations defined in section 2. We will get as immediate consequences the dilation theorems for groups and Gabor type unitary systems introduced in [HL].

In the last section we consider the case of wavelets obtained from a multiresolution analysis. It is known (see [Dau92]) that, unless some restrictions are imposed on the low-pass filter that starts the MRA construction, the wavelets obtained do not form an orthonormal basis but a normalized tight frame. Since such frames can be dilated to orthonormal bases, a natural question would be if the dilation preserves the multiresolution structure. The answer is affirmative and it is given in theorem 5.2 and more concretely in theorem 5.3. In this way we obtain "wavelets" in a Hilbert space bigger than $L^2(\mathbb{R})$.

2. POSITIVE DEFINITE MAPS AND REPRESENTATIONS

We begin this section with a general result of Kolmogorov ([EvLe], [EvKa]). Then we consider several structures and show how to obtain representations from this general theorem.

Definition 2.1. Let X be a nonempty set. We say that a map $K : X \times X \rightarrow \mathbb{C}$ is positive definite, and denote this by $0 \leq K$, if

$$\sum_{i,j=1}^n K(x_i, x_j) \xi_i \bar{\xi}_j \geq 0, \quad (n \in \mathbb{N}, x_i \in X, \xi_i \in \mathbb{C} \text{ for all } i \in \{1, \dots, n\}).$$

Theorem 2.2. [Kolmogorov's theorem] *If $K : X \times X \rightarrow \mathbb{C}$ is positive definite then there exists a Hilbert space H_K and a map $v_K : X \rightarrow H_K$ such that the linear span of $\{v_K(x) \mid x \in X\}$ is dense in H_K and*

$$\langle v_K(x) \mid v_K(y) \rangle = K(x, y), \quad (x, y \in X).$$

Moreover, H_K and v_K are unique up to unitary isomorphisms.

Remark 2.3. Kolmogorov's theorem is valid also for operator-valued positive definite maps and in this form it can be applied for the Stinespring construction and the Naimark-Sz.Nagy dilation. For details consult [EvLe] and [EvKa].

In this paper, for the application to wavelets and Gabor frames, we will need only the more particular version of Kolmogorov's theorem that we mentioned before.

Definition 2.4. If $K : X \times X \rightarrow \mathbb{C}$ is positive definite then we call $[H_K, v_K]$ the representation associated to K .

We note that Kolmogorov's theorem is purely set theoretic; there is no structure on X . We expect that, if X has some additional structure on it and if we assume some compatibility between the positive definite map K and this structure, then the representation associated to K will also be in agreement with the structure of X . In the next examples we will see that this is indeed the case and we review the technique in the case of C^* -algebras and groups.

Example 2.5. [C^* -algebras and the GNS construction] We consider now the case when $X = \mathcal{A}$ is a C^* -algebra and prove that we can obtain the well known GNS construction from Kolmogorov's theorem.

Theorem 2.6. [The GNS construction] *If $\mathcal{A}$ is a C^* -algebra and φ is a positive linear functional on $\mathcal{A}$, then there exists a representation π of $\mathcal{A}$ on a Hilbert space H , that has a cyclic vector $\xi_0 \in H$ such that*

$$\langle \pi(x)\xi_0 | \xi_0 \rangle = \varphi(x), \quad (x \in \mathcal{A}).$$

Proof. The idea is to define $K : \mathcal{A} \times \mathcal{A} \rightarrow \mathbb{C}$ by

$$K(x, y) = \varphi(y^*x), \quad (x, y \in \mathcal{A}).$$

We can use Kolmogorov's theorem to obtain the Hilbert space H_K and the map $v_K : \mathcal{A} \rightarrow H_K$.

For a fixed $x \in \mathcal{A}$, define the operator $\pi(x)$ as follows:

$$\pi(x)(v_K(y)) = v_K(xy), \quad (y \in \mathcal{A}),$$

and extend by linearity. Then everything checks out. $\square$

Example 2.7. [Groups and unitary representations] Take $X = G$ a group. We call $K : G \times G \rightarrow \mathbb{C}$ a group positive definite map if $0 \leq K$ and

$$K(x, y) = K(zx, zy), \quad (x, y, z \in G).$$

We note that such a positive definite map K is uniquely determined by its restriction $\phi(x) = K(x, 1)$ and ϕ is a function of positive type (see [Fol]). The proof of theorem 2.8 will show how the well known correspondence between functions of positive type and unitary representations of groups can be regarded as a consequence of Kolmogorov's theorem.

Theorem 2.8. *Let G be a group and K a group positive definite map on G . Then there exists a unitary representation π_K of G on a Hilbert space H_K with a cyclic vector $\xi_0 \in H_K$ such that*

$$\langle \pi_K(x)\xi_0 | \pi_K(y)\xi_0 \rangle = K(x, y), \quad (x, y \in G).$$

Proof. The proof works exactly as in the case of C^* -algebras: consider $[H_K, v_K]$ the representation associated to K by Kolmogorov's theorem. Define the operators $\pi_K(x)$ for $x \in G$ as follows:

$$\pi_K(x)(v_K(y)) = v_K(xy), \quad (x, y \in G)$$

and extend by linearity. $\square$

Remark 2.9. Note that in the proof of theorem 2.8 we used the representation associated to K and we see that, when K is a group positive definite map, this representation has the unitary representation π_K attached to it. The same observation can be done for the GNS construction: the representation of the C^* -algebra is attached to the representation v_K . This confirms our expectation: when the positive definite map has some compatibility with the existent structure on X , this compatibility projects a nice structure on the associated representation $[H_K, v_K]$. This is the idea that we use throughout this section.

Example 2.10. [Gabor type unitary systems] We recall that a Gabor system is associated to two positive constants $a, b > 0$ and a function $g \in L^2(\mathbb{R})$ and is defined by

$$g_{m,n}(\xi) = e^{2\pi i m b \xi} g(\xi - na), \quad (\xi \in \mathbb{R}).$$

The Gabor systems are one of the major subjects in the study of frames and wavelet theory. If we define the unitary operators U, V on $L^2(\mathbb{R})$,

$$(Uf)(\xi) = e^{2\pi i b \xi} f(\xi), \quad (f \in L^2(\mathbb{R})),$$

$$(Vf)(\xi) = f(\xi - a), \quad (f \in L^2(\mathbb{R})),$$

then $g_{m,n} = U^m V^n g$, $(m, n \in \mathbb{Z})$, and U and V satisfy the relation

$$UV = e^{2\pi i ab} VU.$$

Following [HL], if U and V are unitary operators on a Hilbert space H that verify the relation

$$UV = \lambda VU$$

for some unimodular scalar λ , we then call $\{U^m V^n \mid m, n \in \mathbb{Z}\}$ a Gabor type unitary system. We will prove that these systems fit into our general framework and we construct representations for them.

Theorem 2.11. *Suppose λ is a unimodular scalar and $K : \mathbb{Z}^2 \times \mathbb{Z}^2 \rightarrow \mathbb{C}$ is a positive definite map satisfying*

$$(2.1) \quad K((m+1, n), (m'+1, n')) = K((m, n), (m', n')), \quad (m, n, m', n' \in \mathbb{Z}),$$

$$(2.2) \quad K((m, n+1), (m', n'+1)) = \lambda^{m-m'} K((m, n), (m', n')), \quad (m, n, m', n' \in \mathbb{Z}).$$

Then, on the Hilbert space H_K , there are unitaries U, V and a vector $\xi_0 \in H_K$ such that

$$(2.3) \quad UV = \lambda VU$$

$$(2.4) \quad \{U^m V^n \xi_0 \mid m, n \in \mathbb{Z}\} \text{ is dense in } H_K$$

$$(2.5) \quad \langle U^m V^n \xi_0 \mid U^{m'} V^{n'} \xi_0 \rangle = K((m, n), (m', n')), \quad (m, n, m', n' \in \mathbb{Z}).$$

Moreover, this representation is unique up to unitary isomorphism.

Proof. Let $v_K : \mathbb{Z}^2 \rightarrow H_K$ be the representation associated to K . Define the operators U and V as follows:

$$U(v_K(m, n)) = v_K(m+1, n), \quad (m, n \in \mathbb{Z}),$$

$$V(v_K(m, n)) = \lambda^{-m} v_K(m, n+1), \quad (m, n \in \mathbb{Z}),$$

and then extend by linearity. We check that U, V are well defined and isometric. Take $a_i \in \mathbb{C}$, $(m_i, n_i) \in \mathbb{Z}^2$, $(i \in \{1, \dots, p\})$.

$$\begin{aligned} & \left\langle V\left(\sum_{i=1}^p a_i v_K(m_i, n_i)\right) \mid V\left(\sum_{i=1}^p a_i v_K(m_i, n_i)\right) \right\rangle = \\ &= \left\langle \sum_{i=1}^p a_i \lambda^{-m_i} v_K(m_i, n_i+1) \mid \sum_{i=1}^p a_i \lambda^{-m_i} v_K(m_i, n_i+1) \right\rangle \\ &= \sum_{i,j=1}^p a_i \bar{a}_j \lambda^{-m_i} \overline{\lambda^{-m_j}} K((m_i, n_i+1), (m_j, n_j+1)) \\ &= \sum_{i,j=1}^p a_i \bar{a}_j K((m_i, n_i), (m_j, n_j)) = \left\langle \sum_{i=1}^p a_i v_K(m_i, n_i) \mid \sum_{i=1}^p a_i v_K(m_i, n_i) \right\rangle. \end{aligned}$$

A similar calculation shows that U is well defined and isometric. Since the linear span of the vectors $v_K(m, n)$ is dense in H_K , we can extend U and V to unitaries on H_K .

Next, we check (2.3). Take $(m, n) \in \mathbb{Z}^2$.

$$\begin{aligned} UV v_K(m, n) &= U(\lambda^{-m} v_K(m, n+1)) = \lambda^{-m} v_K(m+1, n+1) \\ &= \lambda V(v_K(m+1, n)) = \lambda VU(v_K(m, n)), \end{aligned}$$

and (2.3) follows by density. Also, note that, if $\xi_0 = v_K(0, 0)$, then

$$U^m V^n \xi_0 = U^m V^n v_K(0, 0) = U^m v_K(0, n) = v_K(m, n), \quad (m, n \in \mathbb{Z}).$$

This will imply (2.4) and (2.5). The uniqueness is a consequence of the uniqueness part of Kolmogorov's theorem. $\square$

Remark 2.12. Any Gabor type unitary system U, V on a Hilbert space H , that has a vector $\xi_0 \in H$ with the property that the linear span of

$$\{U^m V^n \xi_0 \mid m, n \in \mathbb{Z}\}$$

is dense in H , gives rise to a positive definite map K on $\mathbb{Z}^2$ that satisfies (2.1), (2.2) as follows:

$$K((m, n), (m', n')) = \langle U^m V^n \xi_0 \mid U^{m'} V^{n'} \xi_0 \rangle, \quad (m, n, m', n' \in \mathbb{Z}).$$

(2.1), (2.2) are just immediate consequences of the fact that U and V are unitary and $UV = \lambda VU$.

Example 2.13. [Wavelet representations] We recall briefly some facts about wavelet representations. Wavelet theory deals with two unitary operators U and T on $L^2(\mathbb{R})$, corresponding to the integer $N \geq 2$ called the scale:

$$Uf(x) = \frac{1}{\sqrt{N}} f\left(\frac{x}{N}\right), \quad Tf(x) = f(x-1), \quad (x \in \mathbb{R}, f \in L^2(\mathbb{R})).$$

A wavelet is a function $\psi \in L^2(\mathbb{R})$ such that

$$\{U^m T^n \psi \mid m, n \in \mathbb{Z}\}$$

is an orthonormal basis for $L^2(\mathbb{R})$. One way to construct wavelets is by multiresolutions and scaling functions (see [Dau92]). Scaling functions satisfy equations of the form

$$(2.6) \quad U\varphi = \sum_{k \in \mathbb{Z}} a_k T^k \varphi,$$

where a_k are complex coefficients.

The scaling equation can be reformulated using representations. There is a representation of $L^\infty(\mathbb{T})$ ($\mathbb{T}$ is the unit circle) on $L^2(\mathbb{R})$ given by

$$(\widehat{\pi(f)\xi}) = f\widehat{\xi}, \quad (f \in L^\infty(\mathbb{R}), \xi \in L^2(\mathbb{R}))$$

($\widehat{\xi}$ denotes the Fourier transform of ξ and functions on $\mathbb{T}$ are identified with 2π -periodic functions on $\mathbb{R}$). Using this representation, (2.6) can be rewritten as

$$U\varphi = \pi(m_0)\varphi,$$

$m_0(e^{-i\theta}) = \sum_{k \in \mathbb{Z}} a_k e^{-ik\theta}$ is called a low-pass filter. Also, the representation satisfies

$$U\pi(f)U^{-1} = \pi(f(z^N)), \quad (f \in L^\infty(\mathbb{T})).$$

$(U, \pi, L^2(\mathbb{R}), \varphi)$ is called the wavelet representation with scaling function φ .

The wavelet theory has shown a strong interconnection between properties of the scaling function φ and spectral properties of the transfer operator associated to the low-pass filter m_0 :

$$R_{m_0, m_0} f(z) = \frac{1}{N} \sum_{w^N = z} |m_0|^2(w) f(w), \quad (z \in \mathbb{T}, f \in L^1(\mathbb{T})).$$

where $\mathbb{T}$ is endowed with the normalized Haar measure. For more information on this we refer the reader to [BraJo]. In particular, functions that are harmonic with respect to R_{m_0, m_0} , i.e. $R_{m_0, m_0} h = h$, play an important role in the theory.

We recall here a theorem from [Jor98] which establishes the link between functions which are harmonic with respect to R_{m_0, m_0} and wavelet representations, because it is another particularized instance of Kolmogorov's theorem.

Theorem 2.14. *If $m_0 \in L^\infty(\mathbb{T})$ is non-singular (i.e. it doesn't vanish on a set of positive measure) and $h \in L^1(\mathbb{T})$, satisfies*

$$R_{m_0, m_0} h = h, \quad h \geq 0,$$

then there exists a Hilbert space H_h , a representation π_h of $L^\infty(\mathbb{T})$ on H_h , a unitary U_h on H_h and a vector $\varphi_h \in H_h$ such that

$$\overline{\text{span}}\{U_h^{-n} \pi_h(f) \varphi_h \mid n \in \mathbb{N}, f \in L^\infty(\mathbb{T})\} = H_h;$$

$$U_h \pi_h(f) U_h^{-1} = \pi_h(f(z^N)), \quad (f \in L^\infty(\mathbb{T}));$$

$$U_h \varphi_h = \pi_h(m_0) \varphi_h;$$

$$\langle \pi_h(f) \varphi_h | \varphi_h \rangle = \int_{\mathbb{T}} f h d\mu.$$

Moreover, this is unique up to unitary equivalence.

Proof. We give here only a sketch of the proof that uses Kolmogorov's theorem, the rest are calculations which can be found in [Jor98].

Let

$$X = \{(f, n) \mid f \in L^\infty(\mathbb{T}), n \in \mathbb{N}\}.$$

We want to define a positive definite map K on X such that in the end $v_K(f, n) = U_h^{-n} \pi_h(f) \varphi_h$. Then, we must have

$$\begin{aligned} K((f, n), (g, m)) &= \langle U_h^{-n} \pi_h(f) \varphi_h \mid U_h^{-m} \pi_h(g) \varphi_h \rangle \\ &= \langle U_h^m \pi_h(f) \varphi_h \mid U_h^n \pi_h(g) \varphi_h \rangle \\ &= \left\langle \pi_h(f(z^{N^m}) m_0^{(m)}(z)) \varphi_h \mid \pi_h(g(z^{N^n}) m_0^{(n)}(z)) \varphi_h \right\rangle \\ &= \int_{\mathbb{T}} f(z^{N^m}) m_0^{(m)}(z) \overline{g(z^{N^n}) m_0^{(n)}(z)} h d\mu, \end{aligned}$$

where $m_0^{(m)}(z) = m_0(z) m_0(z^N) \dots m_0(z^{N^{m-1}})$.

So we have to define, for $(f, n), (g, m) \in X$,

$$K((f, n), (g, m)) = \int_{\mathbb{T}} f(z^{N^m}) m_0^{(m)}(z) \overline{g(z^{N^n}) m_0^{(n)}(z)} h d\mu.$$

K can be checked to be positive definite so it induces a representation (H_h, v_h) , according to Kolmogorov's theorem. Then, define $\varphi_h = v_h(1, 0)$,

$$U_h v_h(f, 0) = v_h(f(z^N) m_0, 0), \quad (f \in L^\infty(\mathbb{T})),$$

$$U_h v_h(f, n) = v_h(f, n-1), \quad (n \geq 1, f \in L^\infty(\mathbb{T})),$$

and extend by linearity and density.

$$\pi_h(f) v_h(g, n) = v_h(f(z^{N^n}) g(z), n), \quad (f, g \in L^\infty(\mathbb{T}), n \in \mathbb{N}),$$

and extend by linearity and density.

Everything can be checked out as the reader may see in [Jor98]. $\square$

3. INTERTWINING OPERATORS

In the previous section we saw how positive definite maps induce representations on Hilbert spaces. Now we will show that intertwining operators can be constructed in a similar way from maps $L : X \times X \rightarrow \mathbb{C}$ which satisfy some boundedness condition. We will also see that, when X has some structure on it and L is compatible with this structure, then the intertwining operator induced by L will be compatible with the extra structure existent on the induced representations, i.e. the operator is indeed intertwining.

The format of this section is similar to the format of the previous one. We begin with a general, set theoretic result and then particularize it to various structures to obtain more information.

Definition 3.1. Consider two positive definite maps $K, K' : X \times X \rightarrow \mathbb{C}$ and $L : X \times X \rightarrow \mathbb{C}$ (not necessarily positive definite). We say that L is bounded with respect to K and K' if there is a constant $c > 0$ such that

$$(3.1) \quad \left| \sum_{i=1}^m \sum_{j=1}^n L(x_i, y_j) \xi_i \bar{\eta}_j \right|^2 \leq c \left(\sum_{i,i'=1}^m K(x_i, x_{i'}) \xi_i \bar{\xi}_{i'} \right) \left(\sum_{j,j'=1}^n K'(y_j, y_{j'}) \eta_j \bar{\eta}_{j'} \right)$$

for all $x_i, y_j \in X$, $\xi_i, \eta_j \in \mathbb{C}$, $i \in \{1, \dots, m\}, j \in \{1, \dots, n\}$. We denote this by

$$L^2 \leq c K K'.$$

Theorem 3.2. Suppose X is a nonempty set and K, K' are positive definite maps on X . If $L : X \times X \rightarrow \mathbb{C}$ and $L^2 \leq cKK'$ for some $c > 0$, then there exists a unique bounded linear operator $S : H_K \rightarrow H_{K'}$ such that

$$(3.2) \quad \langle Sv_K(x) | v_{K'}(y) \rangle = L(x, y), \quad (x, y \in X).$$

($(H_K, v_K), (H_{K'}, v_{K'})$ are the representation induced by K and K' respectively, according to Kolmogorov's theorem). Moreover, $\|S\| \leq \sqrt{c}$. Conversely, if $S : H_K \rightarrow H_{K'}$ is a bounded linear operator, then there is a unique map $L : X \times X \rightarrow \mathbb{C}$ with $L^2 \leq \|S\|^2 KK'$ that satisfies (3.2).

Proof. Define $B : H_K \times H_{K'} \rightarrow \mathbb{C}$ as follows: for $x_i, y_j \in X, \xi_i, \eta_j \in \mathbb{C}$,

$$B\left(\sum_{i=1}^n \xi_i v_K(x_i), \sum_{j=1}^n \eta_j v_{K'}(y_j)\right) = \sum_{i=1}^m \sum_{j=1}^n \xi_i \bar{\eta}_j L(x_i, y_j).$$

Because $L^2 \leq cKK'$, we have

$$\begin{aligned} \left| B\left(\sum_{i=1}^n \xi_i v_K(x_i), \sum_{j=1}^n \eta_j v_{K'}(y_j)\right) \right|^2 &\leq c \left(\sum_{i,i'=1}^m K(x_i, x_{i'}) \xi_i \bar{\xi}_{i'} \right) \\ &\cdot \left(\sum_{j,j'=1}^n K'(y_j, y_{j'}) \eta_j \bar{\eta}_{j'} \right) = c \left\| \sum_{i=1}^n \xi_i v_K(x_i) \right\|_{H_K}^2 \left\| \sum_{j=1}^n \eta_j v_{K'}(y_j) \right\|_{H_{K'}}^2 \end{aligned}$$

This shows that B is a well defined bounded sesquilinear map which can be extended (by the density properties of v_K and $v_{K'}$) to a bounded sesquilinear map $B : H_K \times H_{K'} \rightarrow \mathbb{C}$. Then there exists a bounded linear operator $S : H_K \rightarrow H_{K'}$ such that $\|S\| \leq \sqrt{c}$ and

$$B(v_1, v_2) = \langle Sv_1 | v_2 \rangle, \quad (v_1 \in H_K, v_2 \in H_{K'}).$$

In particular, one obtains (3.2).

The uniqueness is clear because the spans of $\{v_K(x) | x \in X\}$ and $\{v_{K'}(y) | y \in X\}$ are dense. The converse is also easy, one needs to check that the map L defined by (3.2) satisfies $L^2 \leq \|S\|^2 KK'$, but this is a consequence of Schwarz's inequality. $\square$

Definition 3.3. We call the operator S associated to L in theorem 3.2, the intertwining operator associated to L .

We will also be interested in subrepresentations and in the commutant of a representation. In these instances we will work with only one positive definite map K . We give here a definition which will be appropriate for these situations.

Definition 3.4. Consider K, K' , two positive definite maps on a nonempty set X and a constant $c > 0$. We denote

$$K' \leq cK$$

if, for all $x_i \in X$ and $\xi_i \in \mathbb{C}$, ($i \in \{1, \dots, n\}$),

$$\sum_{i,j=1}^n K'(x_i, x_j) \xi_i \bar{\xi}_j \leq c \sum_{i,j=1}^n K(x_i, x_j) \xi_i \bar{\xi}_j.$$

Proposition 3.5. If K and K' are positive definite maps and $c > 0$ then $K' \leq cK$ if and only if $K'^2 \leq c^2 KK$.

Proof. Suppose $K' \leq cK$. Take $x_i, y_j \in X, \xi_i, \eta_j \in \mathbb{C}$.

$$\begin{aligned} \left| \sum_{i=1}^m \sum_{j=1}^n K'(x_i, y_j) \xi_i \bar{\eta}_j \right|^2 &= \left| \left\langle \sum_{i=1}^m \xi_i v_{K'}(x_i) \mid \sum_{j=1}^n \eta_j v_{K'}(y_j) \right\rangle \right|^2 \\ &\leq \left\| \sum_{i=1}^m \xi_i v_{K'}(x_i) \right\|_{H_{K'}}^2 \left\| \sum_{j=1}^n \eta_j v_{K'}(y_j) \right\|_{H_{K'}}^2 \end{aligned}$$

$$\begin{aligned}
&= \left(\sum_{i,i'=1}^m K'(x_i, x_{i'}) \xi_i \bar{\xi}_{i'} \right) \left(\sum_{j,j'=1}^m K'(y_j, y_{j'}) \eta_j \bar{\eta}_{j'} \right) \\
&\leq c^2 \left(\sum_{i,i'=1}^m K(x_i, x_{i'}) \xi_i \bar{\xi}_{i'} \right) \left(\sum_{j,j'=1}^m K(y_j, y_{j'}) \eta_j \bar{\eta}_{j'} \right).
\end{aligned}$$

Hence $K'^2 \leq c^2 KK$. Conversely, if $K'^2 \leq c^2 KK$ then just take $m = n$, $x_i = y_i$, $\xi_i = \eta_i$ in (3.1) to obtain exactly $K' \leq cK$. $\square$

Corollary 3.6. *Suppose K is positive definite on X . Then, for every positive definite map K' with $K' \leq cK$ for some $c > 0$, there exists a unique positive operator $S : H_K \rightarrow H_K$ with*

$$(3.3) \quad \langle Sv_K(x) | v_K(y) \rangle = K'(x, y), \quad (x, y \in X).$$

Moreover, $\|S\| \leq c$. Conversely, for every positive operator S on H_K there is a unique positive definite map on X that satisfies (3.3). In addition, $K' \leq \|S\|K$.

Proof. Using proposition 3.5 and theorem 3.2, we find an operator S on H_K that satisfies (3.3) and $\|S\| \leq c$. S is positive because

$$\left\langle S \left(\sum_{i=1}^n \xi_i v_K(x_i) \right) \middle| \sum_{i=1}^n \xi_i v_K(x_i) \right\rangle = \sum_{i,j=1}^n \xi_i \bar{\xi}_j K'(x_i, x_j) \geq 0$$

and $\{v_K(x) | x \in X\}$ span a dense subspace of H_K .

For the converse, when S is given, theorem 3.2 shows that there is a K' satisfying (3.3) and $K'^2 \leq \|S\|^2 KK$. K' is positive because S is, and proposition 3.5 implies $K' \leq \|S\|K$. $\square$

In the remainder of this section we apply theorem 3.2 to the situations when X has some additional structure on it and see how the intertwining operators are in compliance with the extra structure of the representations.

Example 3.7. [C^* -algebras] Consider now $X = \mathcal{A}$, a C^* -algebra. We saw in example 2.5 that, when the positive definite map $K : \mathcal{A} \times \mathcal{A} \rightarrow \mathbb{C}$ is given by a positive functional $\varphi : \mathcal{A} \rightarrow \mathbb{C}$,

$$K(x, y) = \varphi(y^*x), \quad (x, y \in \mathcal{A}),$$

then the representation induced by K has the GNS construction attached to it. We want to see for what functions $L : \mathcal{A} \times \mathcal{A} \rightarrow \mathbb{C}$ the associated intertwining operator will intertwine the GNS representations.

Theorem 3.8. *Let $\mathcal{A}$ be a C^* -algebra and φ, φ' two positive functionals on $\mathcal{A}$. Suppose that $\varphi_0 : \mathcal{A} \rightarrow \mathbb{C}$ is linear and $\varphi_0^2 \leq c\varphi\varphi'$ for some $c > 0$ in the sense that*

$$(3.4) \quad |\varphi_0(y^*x)|^2 \leq c\varphi(x^*x)\varphi'(y^*y), \quad (x, y \in \mathcal{A}).$$

Then there exists a unique bounded operator $S : H_\varphi \rightarrow H_{\varphi'}$ such that

$$(3.5) \quad S\pi_\varphi(x) = \pi_{\varphi'}(x)S, \quad (x \in \mathcal{A}),$$

$$(3.6) \quad \langle S\pi_\varphi(x)\xi_0 | \pi_{\varphi'}(y)\xi'_0 \rangle = \varphi_0(y^*x), \quad (x, y \in \mathcal{A}).$$

(Here $(H_\varphi, \pi_\varphi, \xi_0)$ and $(H_{\varphi'}, \pi_{\varphi'}, \xi'_0)$ are the GNS representations associated to φ and φ' respectively (see theorem 2.6).) Moreover $\|S\| \leq \sqrt{c}$. Conversely, if $S : H_\varphi \rightarrow H_{\varphi'}$ is a bounded operator that satisfies (3.5) then there is a unique linear map $\varphi_0 : \mathcal{A} \rightarrow \mathbb{C}$ that satisfies (3.6). In addition (3.4) holds with $c = \|S\|^2$.

Proof. Let $K_\varphi, K_{\varphi'} : \mathcal{A} \times \mathcal{A} \rightarrow \mathbb{C}$,

$$K_\varphi(x, y) = \varphi(y^*x), K_{\varphi'}(x, y) = \varphi'(y^*x), \quad (x, y \in \mathcal{A}).$$

Recall that $H_\varphi = H_{K_\varphi}, H_{\varphi'} = H_{K_{\varphi'}}, \pi_\varphi(x)\xi_0 = v_{K_\varphi}(x), \pi_{\varphi'}(x)\xi'_0 = v_{K_{\varphi'}}(x)$ (see the proof of theorem 2.6).

Define $L(x, y) = \varphi_0(y^*x)$ for $x, y \in \mathcal{A}$. Then (3.4) implies $L^2 \leq cK_\varphi K_{\varphi'}$. Theorem 3.2 gives an operator S with

$$\langle Sv_{K_\varphi}(x) | v_{K_{\varphi'}}(y) \rangle = L(x, y), \quad (x, y \in \mathcal{A});$$

then one checks that S satisfies all the requirements. $\square$

As a corollary we deduce a basic fact about positive operators in the commutant of the GNS representation (see[BraRo]).

Corollary 3.9. *Let φ, φ' be two positive functionals on a C^* -algebra $\mathcal{A}$, $\varphi' \leq c\varphi$ for some $c > 0$ (i.e. $\varphi'(x) \leq c\varphi(x)$ for all positive $x \in \mathcal{A}$). There exists a unique positive linear operator S in the commutant of the GNS representation corresponding to φ such that*

$$(3.7) \quad \langle S\pi_\varphi(x)\xi_0 | \pi_\varphi(y)\xi_0 \rangle = \varphi'(y^*x), \quad (x, y \in \mathcal{A}).$$

Conversely, for any positive operator S in the commutant of $\pi_\varphi(\mathcal{A})$, there is a unique positive functional φ' on $\mathcal{A}$ such that (3.7) holds and $\varphi' \leq \|S\|\varphi$.

Example 3.10. [Groups] Take now $X = G$ a group. We know from theorem 2.8 that, if $K : G \times G \rightarrow \mathbb{C}$ is positive definite and satisfies

$$K(x, y) = K(zx, zy), \quad (x, y, z \in G),$$

then K induces a unitary representation of G on H_K . In the next theorem we look at operators that intertwine these representations.

Theorem 3.11. *Suppose G is a group and K, K' are positive definite maps on G satisfying*

$$K(x, y) = K(zx, zy), \quad K'(x, y) = K'(zx, zy), \quad (x, y, z \in G).$$

Let $L : G \times G \rightarrow \mathbb{C}$ with $L^2 \leq cKK'$ for some $c > 0$. If

$$(3.8) \quad L(x, y) = L(zx, zy), \quad (x, y, z \in G),$$

then there is a unique operator $S : H_K \rightarrow H_{K'}$ such that

$$(3.9) \quad S\pi_K(x) = \pi_{K'}(x)S, \quad (x \in G),$$

$$(3.10) \quad \langle S\pi_K(x)\xi_0 | \pi_{K'}(y)\xi'_0 \rangle = L(x, y), \quad (x \in G).$$

((H_K, π_K, ξ_0), ($H_{K'}, \pi_{K'}, \xi'_0$)) are the unitary representations of G associated to K and K' respectively (see theorem 2.8)). Moreover $\|S\| \leq \sqrt{c}$. Conversely, if $S : H_K \rightarrow H_{K'}$ satisfies (3.9), then there is a unique L that satisfies (3.10). In addition L satisfies (3.8) and $L^2 \leq \|S\|^2 KK'$.

Proof. Recall that, if $v_K : G \rightarrow H_K$ and $v_{K'} : G \rightarrow H_{K'}$ are the representations associated to K by Kolmogorov's theorem, then

$$\pi_K(x)\xi_0 = v_K(x), \quad \pi_{K'}(x)\xi'_0 = v_{K'}(x), \quad (x \in G)$$

(see the proof of theorem 2.8).

Theorem 3.2 implies the existence of an operator $S : H_K \rightarrow H_{K'}$ with $\|S\| \leq \sqrt{c}$ and

$$\langle Sv_K(x) | v_{K'}(y) \rangle = L(x, y), \quad (x, y \in G).$$

The rest follows. $\square$

Corollary 3.12. *Let K, K' be two positive definite maps on the group G that satisfy*

$$K(x, y) = K(zx, zy), \quad K'(x, y) = K'(zx, zy), \quad (x, y, z \in G),$$

and $K' \leq cK$ for some $c > 0$. Then there exists a unique positive operator S on H_K in the commutant of the unitary representation $\pi_K(G)$, such that

$$(3.11) \quad \langle S\pi_K(x)\xi_0 | \pi_K(y)\xi_0 \rangle = K'(x, y), \quad (x, y \in G).$$

Conversely, for every positive operator S in the commutant of $\pi_K(G)$, there is a unique positive definite map K' on G that satisfies (3.11) and

$$K'(x, y) = K'(zx, zy), \quad (x, y, z \in G).$$

Proof. It is an immediate consequence of theorem 3.11. It can also be proved from corollary 3.6. $\square$

Example 3.13. [Gabor type unitary systems] We proved in theorem 2.11 that, given a unimodular $\lambda \in \mathbb{C}$ and a positive definite map K on $\mathbb{Z}^2$ that satisfies

$$(3.12) \quad K((m+1, n), (m'+1, n')) = K((m, n), (m', n')), \quad (m, n, m', n' \in \mathbb{Z}),$$

$$(3.13) \quad K((m, n+1), (m', n'+1)) = \lambda^{m-m'} K((m, n), (m', n')), \quad (m, n, m', n' \in \mathbb{Z}),$$

there is a Gabor type unitary system on H_K generated by two unitaries U_K and V_K . As the reader probably expects, we look at the operators that intertwine these systems.

Theorem 3.14. *Let $\lambda \in \mathbb{C}$, $|\lambda| = 1$ and K, K' positive definite maps on $\mathbb{Z}^2$ satisfying the corresponding relations (3.12) and (3.13). Let $L : \mathbb{Z}^2 \times \mathbb{Z}^2 \rightarrow \mathbb{C}$ with the property that $L^2 \leq cKK'$ for some $c > 0$. If L satisfies the relations (3.12) and (3.13), (with K replaced by L , of course), then there is a unique operator $S : H_K \rightarrow H_{K'}$ such that*

$$(3.14) \quad SU_K = U_{K'}S, \quad SV_K = V_{K'}S,$$

$$(3.15) \quad \left\langle SU_K^m V_K^n \xi_0 \mid U_{K'}^{m'} V_{K'}^{n'} \xi'_0 \right\rangle = L((m, n), (m', n')), \quad (m, n, m', n' \in \mathbb{Z}).$$

($(U_K, V_K, \xi_0), (U_{K'}, V_{K'}, \xi'_0)$ are given by theorem 2.11)). Moreover $\|S\| \leq \sqrt{c}$. Conversely, if $S : H_K \rightarrow H_{K'}$ satisfies (3.14), then there exists a unique $L : \mathbb{Z}^2 \times \mathbb{Z}^2 \rightarrow \mathbb{C}$ that verifies (3.15) and in addition L will verify (3.12) and (3.13), too and $L^2 \leq \|S\|^2 KK'$.

Proof. We recall that $U_K^m V_K^n \xi_0 = v_K(m, n)$ and similarly for K' , ($m, n \in \mathbb{Z}$) (see the proof of theorem 2.11). Theorem 3.2 shows that there is an operator $S : H_K \rightarrow H_{K'}$ with $\|S\| \leq \sqrt{c}$ and such that (3.15) holds. We need to check (3.14). Take $m, n, m', n' \in \mathbb{Z}$ and compute:

$$\begin{aligned} \left\langle SU_K U_K^m V_K^n \xi_0 \mid U_{K'}^{m'} V_{K'}^{n'} \xi'_0 \right\rangle &= L((m+1, n), (m', n')) \\ \left\langle U_{K'} SU_K^m V_K^n \xi_0 \mid U_{K'}^{m'} V_{K'}^{n'} \xi'_0 \right\rangle &= \left\langle SU_K^m V_K^n \xi_0 \mid U_{K'}^{m'-1} V_{K'}^{n'} \xi'_0 \right\rangle \\ &= L((m, n), (m'-1, n')) = L((m+1, n), (m', n')). \end{aligned}$$

The density of the linear spans of $\{U_K^m V_K^n \xi_0 \mid m, n \in \mathbb{Z}\}$ and $\{U_{K'}^{m'} V_{K'}^{n'} \xi'_0 \mid m', n' \in \mathbb{Z}\}$ implies $SU_K = U_{K'}S$. A similar calculation shows that $SV_K = V_{K'}S$.

The converse follows from theorem 3.2: if L is defined by (3.15), the only thing that remains to be verified is that L satisfies (3.12) and (3.13), but this is a consequence of (3.14) and $U_K V_K = \lambda V_K U_K$, $U_{K'} V_{K'} = \lambda V_{K'} U_{K'}$. $\square$

Corollary 3.15. *If K, K' are positive definite maps on $\mathbb{Z}^2$ satisfying the relations (3.12) and (3.13) and $K' \leq cK$ then there is a unique positive definite operator S on H_K that commutes with U_K and V_K and*

$$(3.16) \quad \left\langle SU_K^m V_K^n \xi_0 \mid U_K^{m'} V_K^{n'} \xi_0 \right\rangle = K'((m, n), (m', n')), \quad (m, n, m', n' \in \mathbb{Z}).$$

Conversely, if S is a positive operator that commutes with U_K and V_K then K' defined by (3.16) satisfies (3.12) and (3.13).

Proof. The proof follows the same lines as before. $\square$

Remark 3.16. Theorem 3.2 gives us a general existence result for intertwining operators. The next theorems answer the question what conditions should be imposed on L to obtain that its associated operator S intertwines the extra structure existent on H_K ? We saw that for C^* -algebras the necessary and sufficient condition is that $L(x, y) = \varphi_0(y^*x)$ for some linear φ_0 , for groups we must have $L(x, y) = L(zx, zy)$ and for Gabor type unitary systems, L must satisfy the relations (3.12) and (3.13).

Example 3.17. [Intertwiners of wavelet representations] We mentioned in example 2.13 and theorem 2.14 how wavelet representations can be associated to positive functions $h \in L^1(\mathbb{T})$ with $R_{m_0, m_0}h = h$. In [Dut1] and [Dut2] we studied the operators that intertwine these representations. We indicate now how these can be connected to Kolmogorov's theorem. So we will recall the results from [Dut1] and we sketch the proof based on theorem 3.2.

Given h as in theorem 2.14 call $(U_h, \pi_h, H_h, \varphi_h)$ the cyclic representation of $\mathfrak{A}_N$ associated to h . Also, define the transfer operator associated to a pair $m_0, m'_0 \in L^\infty(\mathbb{T})$ by

$$R_{m_0, m'_0}f(z) = \frac{1}{N} \sum_{w^N=z} m_0(w) \overline{m'_0(w)} f(w), \quad (z \in \mathbb{T}, f \in L^1(\mathbb{T})).$$

Theorem 3.18. *Let $m_0, m'_0 \in L^\infty(\mathbb{T})$ be non-singular and $h, h' \in L^1(\mathbb{T})$, $h, h' \geq 0$, $R_{m_0, m_0}(h) = h$, $R_{m'_0, m'_0}(h') = h'$. Let (U, π, H, φ) , (U', π', H', φ') be the cyclic representations corresponding to h and h' respectively.*

If $h_0 \in L^1(\mathbb{T})$, $R_{m_0, m'_0}(h_0) = h_0$ and $|h_0|^2 \leq chh'$ for some $c > 0$ then there exists a unique operator $S : H \rightarrow H'$ such that

$$(3.17) \quad SU = U'S, \quad S\pi(f) = \pi'(f)S, \quad (f \in L^\infty(\mathbb{T}))$$

$$(3.18) \quad \langle S\pi(f)\varphi | \varphi' \rangle = \int_{\mathbb{T}} fh_0 d\mu, \quad (f \in L^\infty(\mathbb{T})).$$

Moreover $\|S\| \leq \sqrt{c}$. Conversely, if S is an operator that satisfies (3.17), then there is a unique $h_0 \in L^1(\mathbb{T})$ with $R_{m_0, m'_0}h_0 = h_0$ such that (3.18) holds. Moreover, $|h_0|^2 \leq \|S\|^2 hh'$.

Proof. Define X as in the proof of theorem 2.14. For all $(f, n), (g, m) \in X$, we want to obtain

$$\begin{aligned} L((f, n), (g, m)) &= \langle SU^{-n}\pi(f)\varphi | U'^{-m}\pi'(g)\varphi' \rangle \\ &= \langle SU^m\pi(f)\varphi | U'^n\pi'(g)\varphi' \rangle \\ &= \left\langle S\pi(f(z^{N^m})m_0^{(m)}(z))\varphi | \pi'(g(z^{N^n})m_0^{(n)}(z))\varphi' \right\rangle \\ &= \int_{\mathbb{T}} f(z^{N^m})m_0^{(m)}(z)\overline{g(z^{N^n})m_0^{(n)}(z)}h_0 d\mu. \end{aligned}$$

Keep the first and the last terms of the equality and this defines L . L will give rise to S by theorem 3.2. For the details of the required computations, see [Dut1]. The converse, can be also obtained from theorem 3.2, but here the generality of theorem 3.2 isn't really needed. $\square$

4. FRAMES AND DILATIONS

Recall that a set $\{x_i | i \in I\}$ of vectors in a Hilbert space H is called a frame if there are two constants $A, B > 0$ such that

$$A\|f\|^2 \leq \sum_{i \in I} |\langle f | x_i \rangle|^2 \leq B\|f\|^2, \quad (f \in H).$$

If $A = B = 1$ the set $\{x_i | i \in I\}$ is called a normalized tight frame.

Frames have been used extensively in applied mathematics for signal processing and data compression. They play a central role in wavelet theory and the analysis of Gabor systems.

In [HL] the normalized tight frames are interpreted as projections of orthonormal bases and it is proved there that Gabor type normalized tight frames can be dilated to Gabor type orthonormal bases, and normalized tight frames generated by groups can be dilated to orthonormal bases generated by the same group (see theorem 3.8 and 4.8 in [HL]). We will revisit these theorems and show that they are immediate consequences of a general result which proves that any normalized tight frame can be dilated to an orthonormal basis in such a way that the extra structure that may exists is preserved under the dilation.

We begin with a proposition that establishes what positive definite maps give rise to normalized tight frames when represented on a Hilbert space.

Proposition 4.1. *Let K be a positive definite map on a set X . Then $\{v_K(x) \mid x \in X\}$ is a normalized tight frame if and only if for all $x_i \in X, \xi_i \in \mathbb{C}, (i \in \{1, \dots, n\})$:*

$$(4.1) \quad \sum_{i,j=1}^n K(x_i, x_j) \xi_i \bar{\xi}_j = \sum_{x \in X} \left| \sum_{i=1}^n K(x_i, x) \xi_i \right|^2.$$

Proof. If $\{v_K(x) \mid x \in X\}$ is a normalized tight frame then take $f = \sum_{i=1}^n \xi_i v_K(x_i)$. The fact that

$$(4.2) \quad \|f\|^2 = \sum_{x \in X} |\langle f \mid v_K(x) \rangle|^2,$$

translates into (4.1). For the converse, we only need to verify (4.2) for f in a dense subset of H_K (see [HeWe] lemma 1.10). Since the linear span of $\{v_K(x) \mid x \in X\}$ is dense in H_K , we can take $f = \sum_{i=1}^n \xi_i v_K(x_i)$ and (4.2) follows from (4.1). $\square$

Definition 4.2. A positive definite map K on X is called a NTF if and only if $\{v_K(x) \mid x \in X\}$ is a normalized tight frame for H_K .

Before we prove our general result we note that, if $\delta : X \times X \rightarrow \mathbb{C}$ is defined by

$$\delta(x, y) = \begin{cases} 1 & , \quad \text{if } x = y \\ 0 & , \quad \text{otherwise} \end{cases}$$

then δ is a positive definite map and $\{v_\delta(x) \mid x \in X\}$ is an orthonormal basis for H_δ .

Proposition 4.3. *If K is a NTF positive definite map on X then $K \leq \delta$.*

Proof. By definition, $\{v_K(x) \mid x \in X\}$ is a normalized tight frame for H_K . Then, by [HL] proposition 1.1, there exists a Hilbert space H containing H_K as a subspace and an orthonormal basis $\{e(x) \mid x \in X\}$ such that, if P is the projection onto H_K , then $Pe(x) = v_K(x)$ for all $x \in X$.

Now take $x_i \in X, \xi_i \in \mathbb{C}, (i \in \{1, \dots, n\})$.

$$\begin{aligned} \sum_{i,j=1}^n K(x_i, x_j) \xi_i \bar{\xi}_j &= \left\langle \sum_{i=1}^n \xi_i v_K(x_i) \mid \sum_{i=1}^n \xi_i v_K(x_i) \right\rangle \\ &= \left\langle P \left(\sum_{i=1}^n \xi_i e(x_i) \right) \mid P \left(\sum_{i=1}^n \xi_i e(x_i) \right) \right\rangle \\ &\leq \|P\| \left\| \sum_{i=1}^n \xi_i e(x_i) \right\|^2 = \sum_{i=1}^n |\xi_i|^2 \\ &= \sum_{i,j=1}^n \delta(x_i, x_j) \xi_i \bar{\xi}_j. \end{aligned}$$

Therefore $K \leq \delta$. $\square$

Theorem 4.4. *If K, K' are NTF positive definite maps on a countable set X , $K \leq cK'$ for some $c > 0$, then there exists an isometry $W : H_K \rightarrow H_{K'}$, W is induced by K , that is*

$$\langle W v_K(x) \mid v_{K'}(y) \rangle = K(x, y), \quad (x, y \in X),$$

the projection P onto WH_K is also induced by K , i.e.

$$\langle P v_{K'}(x) \mid v_{K'}'(y) \rangle = K(x, y), \quad (x, y \in X),$$

and $P v_{K'}(x) = W v_K(x)$.

Proof. Since $K \leq cK'$, by corollary 3.6, there exists a positive operator S on $H_{K'}$ such that

$$\langle S v_{K'}(x) \mid v_{K'}(y) \rangle = K(x, y), \quad (x, y \in X).$$

Since S is positive, it has a positive square root $S^{\frac{1}{2}}$. Then

$$\left\langle S^{\frac{1}{2}} v_{K'}(x) \mid S^{\frac{1}{2}} v_{K'}(y) \right\rangle = \langle S v_{K'}(x) \mid v_{K'}(y) \rangle = K(x, y), \quad (x, y \in X).$$

Take

$$H = \overline{\text{span}}\{S^{\frac{1}{2}}v_{K'}(x) \mid x \in X\}.$$

By the uniqueness part of Kolmogorov's theorem, there is a unitary $W : H_K \rightarrow H$ such that

$$Wv_K(x) = S^{\frac{1}{2}}v_{K'}(x), \quad (x \in X).$$

But then $\{S^{\frac{1}{2}}v_{K'}(x) \mid x \in X\}$ is a normalized tight frame for H . Also, we know that $\{v_{K'}(x) \mid x \in X\}$ is a normalized tight frame for $H_{K'}$. So $S^{\frac{1}{2}} : H_{K'} \rightarrow H$ maps a normalized tight frame to a normalized tight frame, therefore it must be a co-isometry (see [HL] proposition 1.9). It follows that $S^{\frac{1}{2}}(S^{\frac{1}{2}})^* : H \rightarrow H$ is the identity on H so S is the identity on H .

We also know that $\text{range}(S^{\frac{1}{2}}) = \text{range}(S)$. This implies that $S(Sv) = Sv$ for all $v \in H_{K'}$ and, as $S \geq 0$, S is the projection onto H . Consequently, we also have $S = S^{\frac{1}{2}}$ and everything follows now by an easy computation:

$$\begin{aligned} Sv_{K'}(x) &= S^{\frac{1}{2}}v_{K'}(x) = Wv_K(x), \quad (x \in X). \\ \langle Wv_K(x) \mid v_{K'}(y) \rangle &= \langle Sv_{K'}(x) \mid v_{K'}(y) \rangle = K(x, y), \quad (x, y \in X). \end{aligned}$$

□

Remark 4.5. Theorem 4.4 can be used to construct dilation theorems for unitary systems (the reader should have in mind the specific examples of groups and Gabor type unitary systems).

Recall some definitions from [HL]. If $\mathcal{U}$ is a countable set of unitaries on a Hilbert space H , then $\xi \in H$ is called a complete wandering vector (complete normalized tight frame vector) if $\{U\xi \mid U \in \mathcal{U}\}$ is an orthonormal basis (normalized tight frame) for H . A dilation theorem will take the following form:

If $\mathcal{U}$ is a unitary system on a Hilbert space H that has a complete normalized tight frame vector η , then there is a Hilbert space H_1 that contains H and a unitary system $\mathcal{U}_1$ on H_1 such that $\mathcal{U}_1$ has a complete wandering vector ξ and if P is the projection onto H then $P\xi = \eta$, P commutes with $\mathcal{U}_1$ and $U_1 \mapsto U_1|_H$ is an isomorphism of $\mathcal{U}_1$ onto $\mathcal{U}$.

The proof will be guided by the following steps:

1. Construct $K : \mathcal{U} \times \mathcal{U} \rightarrow \mathbb{C}$, $K(x, y) = \langle x\eta \mid y\eta \rangle$; then K is an NTF positive definite map and $H_K = H$, $v_K(x) = x\eta$, ($x \in \mathcal{U}$) and $\mathcal{U}$ is the extra structure $\mathcal{U}_K$ induced by K .
2. Verify that $\delta : \mathcal{U} \times \mathcal{U} \rightarrow \mathbb{C}$ satisfies the required compatibility conditions with $\mathcal{U}$.
3. Construct H_δ , v_δ and the additional structure $\mathcal{U}_\delta$ with cyclic vector ξ_δ which is a complete wandering vector for $\mathcal{U}_\delta$.
4. Since $K \leq \delta$ (proposition 4.3), according to theorem 4.4 there is an isometry $W : H \rightarrow H_\delta$ which is induced by K ; the projection P onto WH is also induced by K and $P\xi_\delta = \eta$. As K is compatible with the structure $\mathcal{U}$, W will intertwine $\mathcal{U}$ and $\mathcal{U}_\delta$ and P commutes with $\mathcal{U}_\delta$. So WH is invariant for $\mathcal{U}_\delta$ and $WUW^{-1} = U_\delta|_H$ for all $U \in \mathcal{U}$ (U_δ is the unitary in $\mathcal{U}_\delta$ that corresponds to U in the representation).
5. Identify H with WH and everything will follow.

We will use the guidelines of remark 4.5 to show how one can obtain the dilation theorems 3.8 and 4.8 from [HL] for groups and Gabor type unitary systems.

Theorem 4.6. [HL] *Suppose $\mathcal{U}$ is a unitary group on H with a complete normalized tight frame vector η . Then there is a Hilbert space H_1 containing H and a unitary group $\mathcal{U}_1$ such that $\mathcal{U}_1$ has a complete wandering vector ξ , if P is the projection onto H then P commutes with $\mathcal{U}_1$, $P\xi = \eta$ and $U_1 \mapsto U_1|_H$ is an isomorphism of $\mathcal{U}_1$ onto $\mathcal{U}$. Consequently, $PU_1\xi = U_1|_H\eta$ for all $U_1 \in \mathcal{U}_1$ (that is the normalized tight frame $\{U\eta \mid U \in \mathcal{U}\}$ can be dilated to the orthonormal basis $\{U_1\xi \mid U_1 \in \mathcal{U}_1\}$).*

Proof. Define $K : \mathcal{U} \times \mathcal{U} \rightarrow \mathbb{C}$, $K(x, y) = \langle x\eta \mid y\eta \rangle$ for $x, y \in \mathcal{U}$. It is clear that K is an NTF positive definite map with

$$(4.3) \quad K(zx, zy) = K(x, y), \quad (x, y, z \in \mathcal{U})$$

and $H_K = H$, $v_K(x) = x\eta$ and the representation π_K given by theorem 2.8 is $\pi_K(x) = x$ for $x \in \mathcal{U}$.

It is also clear that δ satisfies a relation of type (4.3) so it is compatible with the group structure and by theorem 2.8 it induces a cyclic representation $(H_\delta, \pi_\delta, \xi_\delta)$ of $\mathcal{U}$ with $\xi_\delta = v_\delta(1)$ a complete wandering vector.

By proposition 4.3, $K \leq \delta$. By theorem 4.4 there is an isometry $W : H \rightarrow H_\delta$ which is induced by K , the projection P onto WH is also induced by K and $P\xi_\delta = \eta$. Then, by theorem 3.11, W is intertwining that is

$$Wx = \pi(x)W, \quad (x \in \mathcal{U}),$$

and P is in the commutant of $\pi_\delta(\mathcal{U})$. So WH is invariant for all $\pi_\delta(x)$, $x \in \mathcal{U}$ and $WxW^{-1} = \pi_\delta(x)$ for $x \in \mathcal{U}$.

Identify H with WH and define $\mathcal{U}_1 = \pi_\delta(\mathcal{U})$, $\xi = \xi_\delta$ and everything follows. $\square$

Theorem 4.7. [HL] *Let $\mathcal{U} = \{U^m V^n \mid m, n \in \mathbb{Z}\}$ be a Gabor type unitary system associated to λ on a Hilbert space H . Suppose $\mathcal{U}$ has a complete normalized tight frame vector $\eta \in H$. Then there is a Gabor type unitary system $\mathcal{U}_1 (= \{U_1^m V_1^n \mid m, n \in \mathbb{Z}\})$ associated to λ on a Hilbert space H_1 containing H , such that $\mathcal{U}_1$ has a complete wandering vector ξ and if P is the projection onto H then P commutes with U_1 and V_1 , $P\xi = \eta$ and $U = U_1|_H$, $V = V_1|_H$.*

Proof. The proof is analogous to the proof of theorem 4.6, the only difference is to verify that δ satisfies the compatibility relations (3.12) and (3.13) and this is trivial. $\square$

5. A DILATION THEOREM FOR WAVELETS

Let us recall the algorithm for the construction of compactly supported wavelets. For details we refer the reader to [Dau92] for the scale $N = 2$ and to [BraJo97] for arbitrary scale N .

One starts with the low-pass filter $m_0 \in L^2(\mathbb{T})$ which is a trigonometric polynomial that satisfies $m_0(1) = \sqrt{N}$ and the quadrature mirror filter condition

$$(5.1) \quad \frac{1}{N} \sum_{w^N=z} |m_0|^2(w) = 1, \quad (z \in \mathbb{T}).$$

Then define the scaling function $\varphi \in L^2(\mathbb{R})$ by taking the inverse Fourier transform of

$$(5.2) \quad \widehat{\varphi}(x) = \prod_{k=1}^{\infty} \frac{m_0\left(\frac{x}{N^k}\right)}{\sqrt{N}}, \quad (x \in \mathbb{R}).$$

To construct wavelets one needs the high-pass filters $m_1, \dots, m_{N-1} \in L^\infty(\mathbb{T})$ such that the matrix

$$(5.3) \quad \frac{1}{\sqrt{N}} \begin{pmatrix} m_0(z) & m_0(\rho z) & \dots & m_0(\rho^{N-1}z) \\ m_1(z) & m_1(\rho z) & \dots & m_1(\rho^{N-1}z) \\ \vdots & \vdots & \ddots & \vdots \\ m_{N-1}(z) & m_{N-1}(\rho z) & \dots & m_{N-1}(\rho^{N-1}z) \end{pmatrix} \text{ is unitary for a.e } z \in \mathbb{T}.$$

($\rho = e^{\frac{2\pi i}{N}}$).

When $N = 2$ a choice for m_1 can be

$$m_1(z) = z \overline{m_0}(-z) f(z^2), \quad (z \in \mathbb{T}),$$

where $|f(z)| = 1$ on $\mathbb{T}$.

The wavelets are defined as follows:

$$(5.4) \quad \widehat{\psi}_i(x) = \frac{m_i\left(\frac{x}{N}\right)}{\sqrt{N}} \widehat{\varphi}\left(\frac{x}{N}\right), \quad (x \in \mathbb{R}, i \in \{1, \dots, N-1\}),$$

or, in terms of the wavelet representation,

$$(5.5) \quad \psi_i = U^{-1} \pi(m_i) \varphi, \quad (i \in \{1, \dots, N-1\}).$$

It is known that, in order to achieve orthogonality, extra conditions must be imposed on m_0 . If R_{m_0, m_0} has only one continuous fixed point (up to a multiplicative constant), the set

$$\{U^m T^n \psi_i \mid m, n \in \mathbb{Z}, i \in \{1, \dots, N-1\}\}$$

is an orthonormal basis for $L^2(\mathbb{R})$.

However, when this extra condition is not satisfied, one still gets good properties, namely, the fact that the above set is a normalized tight frame for $L^2(\mathbb{R})$. In the sequel, we show how one can dilate this normalized tight frame to an orthonormal basis in such a way that the multiresolution structure is preserved so that "wavelets" in a space bigger than $L^2(\mathbb{R})$ are obtained.

We begin with a proposition that explains the multiresolution structure of the cyclic representations presented in example 2.13.

In the sequel we define $m_0 \in L^\infty(\mathbb{T})$ to be non-singular if the set $\{z \in \mathbb{T} \mid m_0(z) = 0\}$ has zero measure and $|m_0|$ is not constant 1 a.e.

Proposition 5.1. *Let $m_0 \in L^\infty(\mathbb{T})$ be non-singular, $h \in L^1(\mathbb{T})$, $h \geq 0$ and $R_{m_0, m_0} h = h$. Let $(U_h, \pi_h, H_h, \varphi_h)$ be the cyclic representation associated to h . Define $T_h = \pi_h(z)$,*

$$V_0^h = \overline{\text{span}}\{T_h^k \varphi_h \mid k \in \mathbb{Z}\},$$

$$V_j^h = U_h^{-j} V_0^h, \quad (j \in \mathbb{Z}).$$

Then

$$(5.6) \quad U_h T_h U_h^{-1} = T_h^N,$$

$$(5.7) \quad V_j^h \subset V_{j+1}^h, \quad (j \in \mathbb{Z}),$$

$$(5.8) \quad \overline{\bigcup_{j \in \mathbb{Z}} V_j^h} = H_h,$$

$$(5.9) \quad \bigcap_{j \in \mathbb{Z}} V_j^h = \{0\}$$

Assume $h = 1$. Then

$$(5.10) \quad \{T_h^k \varphi_h \mid k \in \mathbb{Z}\} \text{ is an orthonormal basis for } V_0^h.$$

If $m_1, \dots, m_{N-1}$ satisfy (5.3) and

$$(5.11) \quad \psi_i^h = U_h^{-1} \pi_h(m_i) \varphi_h, \quad (i \in \{1, \dots, N-1\}),$$

then

$$(5.12) \quad \{T_h^k \psi_i^h \mid k \in \mathbb{Z}, i \in \{1, \dots, N-1\}\} \text{ is an orthonormal basis for } V_1^h \ominus V_0^h$$

and

$$(5.13) \quad \{U_h^m T_h^n \psi_i^h \mid m, n \in \mathbb{Z}, i \in \{1, \dots, N-1\}\} \text{ is an orthonormal basis for } L^2(\mathbb{R}).$$

Proof. (5.6) follows from $U_h \pi_h(f(z)) U_h^{-1} = \pi_h(f(z^N))$ with $f(z) = z$.

If $f(z) = \sum_{k=-p}^p a_k z^k$ is a trigonometric polynomial then

$$\pi_h(f) \varphi_h = \sum_{k=-p}^p a_k T_h^k \varphi_h \in V_0^h.$$

Each $f \in L^\infty(\mathbb{T})$ is the pointwise limit of a uniformly bounded sequence of trigonometric polynomials, hence, by lemma 2.8 in [Dut1], $\pi_h(f) \varphi_h \in V_0^h$. Then

$$U_h \pi_h(f) \varphi_h = \pi_h(f(z^N)) U_h \varphi_h = \pi_h(f(z^N) m_0(z)) \varphi_h \in V_0^h$$

so $V_{-1}^h \subset V_0^h$ and this implies (5.7).

Also $U_h^{-n} \pi_h(f) \varphi_h \in V_n^h$ for all $f \in L^\infty(\mathbb{T})$ and $n \in \mathbb{Z}$ and (5.8) follows by density. (5.9) is proved in theorem 5.6 from [Jor98].

If $h = 1$ then, for $k \in \mathbb{Z}$,

$$\langle T_h^k \varphi_h \mid \varphi_h \rangle = \langle \pi_h(z^k) \varphi_h \mid \varphi_h \rangle = \int_{\mathbb{T}} z^k d\mu = \delta_{k,0},$$

so (5.10) is valid.

It remains to prove (5.12) because (5.13) follows from this immediately. The argument is essentially the one in [BraJo97] theorem 10.1. We will include it here to make sure everything works.

For $k, l \in \mathbb{Z}$ and $i, j \in \{1, \dots, N-1\}$ we have

$$\begin{aligned} \langle T_h^k \psi_i^h | T_h^l \psi_j^h \rangle &= \langle \pi_h(z^k) U_h^{-1} \pi_h(m_i) \varphi_h | \pi_h(z^l) U_h^{-1} \pi_h(m_j) \varphi_h \rangle \\ &= \langle U_h^{-1} \pi_h(z^{Nk} m_i(z)) \varphi_h | U_h^{-1} \pi_h(z^{Nl} m_j(z)) \varphi_h \rangle \\ &= \int_{\mathbb{T}} z^{N(k-l)} m_i(z) \overline{m_j}(z) d\mu \\ &= \int_{\mathbb{T}} z^{k-l} \frac{1}{N} \sum_{w^N=z} m_i(w) \overline{m_j}(w) d\mu = \delta_{i,j} \delta_{k,l}, \end{aligned}$$

for the last equality we used (5.3). So

$$\{T_h^k \psi_i^h | k \in \mathbb{Z}, i \in \{1, \dots, N-1\}\}$$

is an orthonormal set.

Take, $U_h^{-1} \pi_h(m) \varphi_h \in V_1^h$, $m \in L^\infty(\mathbb{T})$ (vectors of this form are dense in V_1^h). $U_h^{-1} \pi_h(m) \varphi_h \perp V_0^h$ is equivalent to, for all $f \in L^\infty(\mathbb{T})$:

$$\begin{aligned} 0 &= \langle U_h^{-1} \pi_h(m) \varphi_h | \pi_h(f) \varphi_h \rangle = \langle \pi_h(m) \varphi_h | \pi_h(f(z^N) m_0(z)) \varphi_h \rangle \\ &= \int_{\mathbb{T}} m(z) \overline{f}(z^N) \overline{m_0}(z) d\mu \\ &= \int_{\mathbb{T}} \frac{1}{N} \sum_{w^N=z} m(w) \overline{m_0}(w) \overline{f}(z) d\mu \end{aligned}$$

which is equivalent to

$$\frac{1}{N} \sum_{w^N=z} m(w) \overline{m_0}(w) = 0 \text{ a.e. on } \mathbb{T}.$$

This shows in particular that $\psi_i^h \perp V_0^h$ for all $i \in \{1, \dots, N-1\}$. Also, the vector

$$\vec{m}(z) = (m(z), m(\rho z), \dots, m(\rho^{N-1} z))$$

($\rho = e^{-2\pi i/N}$) must be perpendicular to the vector

$$\vec{m}_0(z) = (m_0(z), m_0(\rho z), \dots, m_0(\rho^{N-1} z))$$

for almost all z , so

$$\vec{m}(z) = \sum_{k=1}^{N-1} \mu_k(z) \vec{m}_k(z)$$

where $\mu_k(z) = \langle \vec{m}(z) | \vec{m}_k(z) \rangle$ (which shows that $\mu_k \in L^\infty(\mathbb{T})$).

Since $\vec{m}(\rho z)$ is a circular permutation of $\vec{m}(z)$, it follows that we must have $\mu_k(\rho z) = \mu_k(z)$, that is $\mu_k(z) = \lambda_k(z^N)$ for some $\lambda_k \in L^\infty(\mathbb{T})$.

Then

$$m(z) = \sum_{k=1}^{N-1} \lambda_k(z^N) m_k(z), \quad (z \in \mathbb{T}),$$

and we compute

$$U_h^{-1} \pi_h(m) \varphi_h = U_h^{-1} \pi_h\left(\sum_{k=1}^{N-1} \lambda_k(z^N) m_k(z)\right) \varphi_h = \sum_{k=1}^{N-1} \pi_h(\lambda_k) \psi_k^h.$$

and this shows that

$$U_h^{-1} \pi_h(m) \varphi_h \in \overline{\text{span}}\{T_h^k \psi_i^h | k \in \mathbb{Z}, i \in \{1, \dots, N-1\}\}$$

by an argument similar to the one used in the begining of the proof (now for ψ_i^h instead of φ_h). This completes the proof of (5.12). $\square$

Motivated by the discussion in the begining of this section, we give a dilation theorem for wavelets. The theorem describes how one can dilate a normalized tight frame wavelet to an orthonormal wavelet in a bigger space.

Theorem 5.2. . Let $m_0 \in L^\infty(\mathbb{T})$ be a non-singular filter with $R_{m_0, m_0}1 = 1$, $h \in L^\infty(\mathbb{T})$, $h \geq 0$, $R_{m_0, m_0}h = h$, and consider $(U_h, \pi_h, H_h, \varphi_h)$, the cyclic representation associated to h . Assume also there are given filters $m_1, \dots, m_{N-1} \in L^\infty(\mathbb{T})$ such that (5.3) holds, define ψ_i^h as in (5.11) ($i \in \{1, \dots, N-1\}$) and suppose

$$\{U_h^m T_h^n \psi_i^h \mid m, n \in \mathbb{Z}, i \in \{1, \dots, N-1\}\}$$

is a normalized tight frame for H_h ($T_h = \pi_h(z)$).

Then, if $(U_1, \pi_1, H_1, \varphi_1)$ is the cyclic representation associated to the constant function 1, then there exists an isometry $W : H_h \rightarrow H_1$ with the following properties:

(i) $WU_h = U_1W$, $W\pi_h(f) = \pi_1(f)W$ for all $f \in L^\infty(\mathbb{T})$;

(ii) If P is the projection onto WH_h then

$$(5.14) \quad PU_1 = U_1P, \quad P\pi_1(f) = \pi_1(f)P, \quad (f \in L^\infty(\mathbb{T}));$$

$$(5.15) \quad P\varphi_1 = W\varphi_h.$$

(iii) If $\psi_i^1 = U_1^{-1}\pi_1(m_i)\varphi_1$, $i \in \{1, \dots, N-1\}$ then

$$(5.16) \quad \{U_1^m T_1^n \psi_i^1 \mid m, n \in \mathbb{Z}, i \in \{1, \dots, N-1\}\} \text{ is an orthonormal basis for } H_1,$$

where $T_1 = \pi_1(z)$;

$$(5.17) \quad P\psi_i^1 = W\psi_i^h, \quad (i \in \{1, \dots, N-1\}).$$

Proof. The proof is similar to the one of theorem 4.4 but some additional arguments are needed. Since $h \in L^\infty(\mathbb{T})$, we have $|h|^2 \leq \|h\|_\infty 1$ so, by theorem 3.18 there is a positive operator S on H_1 that commutes with U_1 and π_1 and

$$\langle S\pi_1(f)\varphi_1 \mid \varphi_1 \rangle = \int_{\mathbb{T}} fh d\mu, \quad (f \in L^\infty(\mathbb{T})).$$

S has a positive square root $S^{\frac{1}{2}}$ that commutes with U_1 and π_1 . Also the projection P onto the range H of $S^{\frac{1}{2}}$ must commute with U_1 and π_1 . Then we can restrict π_1 and U_1 to H and

$$\langle \pi_1(f)S^{\frac{1}{2}}\varphi_1 \mid S^{\frac{1}{2}}\varphi_1 \rangle = \int_{\mathbb{T}} fh d\mu,$$

$$U_1 S^{\frac{1}{2}}\varphi_1 = \pi_1(m_0)S^{\frac{1}{2}}\varphi_1.$$

The uniqueness part of theorem 2.14 implies that there is a unitary W from H_h to H with $W\varphi_h = S^{\frac{1}{2}}\varphi_1$, $WU_h = U_1W$ and $W\pi_h(f) = \pi_1(f)W$ for $f \in L^\infty(\mathbb{T})$. From these commuting properties of W and P it follows that

$$W(U_h^m T_h^n \psi_i^h) = S^{\frac{1}{2}}(U_1^m T_1^n \psi_i^1), \quad (m, n \in \mathbb{Z}, i \in \{1, \dots, N-1\}).$$

Hence, $S^{\frac{1}{2}}$ maps an orthonormal basis to a normalized tight frame so it must be a co-isometry. Then, proceeding as in the proof of theorem 4.4 we get that $S = S^{\frac{1}{2}} = P$ and everything follows. $\square$

When m_0 is a regular filter (we will give the precise meaning of that in a moment), we can really get our hands on the abstract cyclic representation associated to the constant function 1 so that we obtain a very concrete dilation theorem for non-orthogonal wavelets in $L^2(\mathbb{R})$. The construction is given in the next theorem and it is based on the results presented in [Dut2].

Before we state the result, some definitions are needed. A vector $(z_1, z_2, \dots, z_p)$ is called an m_0 -cycle if $z_1^N = z_2, z_2^N = z_3, \dots, z_p^N = z_1$, z_i are distinct and $|m_0(z_i)| = \sqrt{N}$ for all $i \in \{1, \dots, p\}$.

For $f \in L^\infty(\mathbb{T})$ and $z_0 \in \mathbb{T}$ define $\alpha_{z_0}(f)(z) = f(z z_0)$ for $z \in \mathbb{T}$. For $n \in \mathbb{N}$

$$m_0^{(n)}(z) = m_0(z)m_0(z^N)\dots m_0(z^{N^{n-1}}), \quad (z \in \mathbb{T}).$$

Theorem 5.3. *Let m_0 be a Lipschitz function on $\mathbb{T}$ with finitely many zeroes $m_0(1) = \sqrt{N}$, $R_{m_0, m_0}1 = 1$. Let $C_j = (z_{1,j}, \dots, z_{p_j,j})$ be the m_0 -cycles, $j \in \{1, \dots, n\}$, $m_0(z_{k,j}) = \sqrt{N}e^{i\theta_{k,j}}$ for all $k \in \{1, \dots, p_j\}$, $j \in \{1, \dots, n\}$, $\theta_j = \theta_{1,j} + \dots + \theta_{p_j,j}$.*

For each $j \in \{1, \dots, n\}$ define: $H_j = L^2(\mathbb{R})^{p_j}$, $U_j : H_j \rightarrow H_j$

$$U_j(\xi_1, \dots, \xi_{p_j}) = \left(e^{i\theta_{1,j}} U \xi_2, \dots, e^{i\theta_{p_j-1,j}} U \xi_{p_j}, e^{i\theta_{p_j,j}} U \xi_1 \right),$$

where $U\xi(x) = \frac{1}{\sqrt{N}}f\left(\frac{x}{N}\right)$ for $\xi \in L^2(\mathbb{R})$.

For $f \in L^\infty(\mathbb{T})$

$$\pi_j(f)(\xi_1, \dots, \xi_{p_j}) = \left(\pi(\alpha_{z_{1,j}}(f))(\xi_1), \dots, \pi(\alpha_{z_{p_j,j}}(f))(\xi_{p_j}) \right),$$

where π is the representation on $L^2(\mathbb{R})$ defined in example 2.13.

$$\widehat{\varphi}_{k,j}(x) = \prod_{l=1}^{\infty} \frac{e^{-i\theta_j} \alpha_{z_{k,j}} \left(m_0^{(p_j)} \right) \left(\frac{x}{N^{lp_j}} \right)}{\sqrt{N^{lp_j}}}, \quad (l \in \{1, \dots, p_j\}),$$

$$\varphi_j = (\varphi_{1,j}, \dots, \varphi_{p_j,j}).$$

Finally, define $H_0 = H_1 \oplus \dots \oplus H_n$, $U_0 = U_1 \oplus \dots \oplus U_n$, $\pi_0(f) = \pi_1(f) \oplus \dots \oplus \pi_n(f)$ for $f \in L^\infty(\mathbb{T})$ and $\varphi_0 = \varphi_1 \oplus \dots \oplus \varphi_n$. Then $(U_0, \pi_0, H_0, \varphi_0)$ is the cyclic representation associated to the constant function 1.

Also if C_1 is the trivial m_0 -cycle $C_1 = (1)$ then $H_1 = L^2(\mathbb{R})$, $U_1 = U$, $\pi_1 = \pi$ and

$$\widehat{\varphi}_1(x) = \prod_{l=1}^{\infty} \frac{m_0\left(\frac{x}{N^l}\right)}{\sqrt{N}}, \quad (x \in \mathbb{R}),$$

so $(U_1, \pi_1, H_1, \varphi_1)$ is the usual wavelet representation on $L^2(\mathbb{R})$.

If $T_j = \pi_j(z)$, for $j \in \{1, \dots, n\}$ then

$$T_j(\xi_1, \dots, \xi_{p_j})(x) = (z_{1,j} T \xi_1, \dots, z_{p_j,j} T \xi_{p_j}),$$

where $T\xi(x) = \xi(x-1)$ for $\xi \in L^2(\mathbb{R})$.

Assume $m_1, \dots, m_{N-1} \in L^\infty(\mathbb{T})$ satisfy (5.3). Define

$$\psi_i^1 = U_1^{-1} \pi_1(m_i) \varphi_1 (\in L^2(\mathbb{R})), \quad \psi_i^0 = U_0^{-1} \pi_0(m_i) \varphi_0, \quad i \in \{1, \dots, N-1\}$$

and let P_1 be the projection from H_0 onto H_1 and $T_0 = T_1 \oplus \dots \oplus T_n$. Then

$$(5.18) \quad P_1 U_0 = U_0 P_1, \quad P_1 T_0 = T_0 P_1, \quad P_1 \pi_0(f) = \pi_0(f) P_1, \quad (f \in L^\infty(\mathbb{T}));$$

$$(5.19) \quad U_0|_{H_1} = U_1 (= U), \quad T_0|_{H_1} = T_1 (= T), \quad \pi_0(f)|_{H_1} = \pi_1(f) (= \pi(f)), \quad (f \in L^\infty(\mathbb{T}));$$

$$(5.20) \quad P_1 \varphi_0 = \varphi_1, \quad P_1 \psi_i^0 = \psi_i^1, \quad (i \in \{1, \dots, N-1\});$$

$$(5.21) \quad U_0 \varphi_0 = \pi_0(m_0) \varphi_0, \quad U_1 \varphi_1 = \pi_1(m_0) \varphi_1;$$

$$(5.22) \quad \{T_0^k \varphi_0 \mid k \in \mathbb{Z}\} \text{ is an orthonormal set};$$

$$(5.23) \quad \{U_0^m T_0^n \psi_i^0 \mid m, n \in \mathbb{Z}, i \in \{1, \dots, N-1\}\} \text{ is an orthonormal basis for } H_0;$$

$$(5.24) \quad \{U_1^m T_1^n \psi_i^1 \mid m, n \in \mathbb{Z}, i \in \{1, \dots, N-1\}\} \text{ is a normalized tight frame for } H_1 = L^2(\mathbb{R}).$$

Proof. The fact that the cyclic representation associated to the constant function 1 is proved in [Dut2], one needs only to take the inverse Fourier transform of the representation presented there to obtain the one described here. Then (5.18) and (5.19) follow trivially from the definition, (5.20) follows from the definition and the commuting properties of P_1 , (5.21) is included in the definition of the cyclic representation, (5.22) and (5.23) are consequences of proposition 5.1 and (5.24) (which is also well known, see [Dau92] or [BraJo97]) follows from the fact that the projection of an orthonormal basis is a normalized tight frame (see [HL]). $\square$

Example 5.4. We apply theorem 5.3 to the low-pass filter

$$m_0(z) = \frac{1+z^3}{\sqrt{2}} = \sqrt{2}e^{-\frac{3i\theta}{2}} \cos\left(\frac{3\theta}{2}\right), \quad (z = e^{-i\theta} \in \mathbb{T})$$

which is known to give non-orthogonal wavelets. The scale $N = 2$. Some short computations show that $m_0(1) = \sqrt{2}$, $R_{m_0, m_0}1 = 1$. The m_0 -cycles are

$$C_1 = (z_{1,1} = 1), C_2 = (z_{2,1} = e^{2\pi i/3}, z_{2,2} = e^{4\pi i/3})$$

$p_1 = 1, p_2 = 2$, $m_0(z_{1,1}) = m_0(z_{2,1}) = m_0(z_{2,2}) = \sqrt{2}$ so $\theta_{1,1} = \theta_{2,1} = \theta_{2,2} = 0$ and $\theta_1 = \theta_2 = 0$.

$$U_0 : L^2(\mathbb{R})^3 \rightarrow L^2(\mathbb{R})^3, \quad U_0(\xi_1, \xi_2, \xi_3) = (U\xi_1, U\xi_3, U\xi_2),$$

$$T_0 : L^2(\mathbb{R})^3 \rightarrow L^2(\mathbb{R})^3, \quad T_0(\xi_1, \xi_2, \xi_3) = (T\xi_1, e^{2\pi i/3}T\xi_2, e^{4\pi i/3}T\xi_3).$$

Then, as $\alpha_{z_{1,1}}(m_0) = m_0$, $\alpha_{z_{2,1}}(m_0) = \alpha_{z_{2,2}}(m_0) = m_0$,

$$\widehat{\varphi}_{1,1}(x) = \prod_{l=1}^{\infty} e^{-\frac{3ix}{2^{l+1}}} \cos\left(\frac{3x}{2^{l+1}}\right) = e^{-\frac{3ix}{2}} \frac{\sin\left(\frac{3x}{2}\right)}{\frac{3x}{2}}, \quad (x \in \mathbb{R}),$$

$$\widehat{\varphi}_{2,1}(x) = \prod_{l=1}^{\infty} \frac{m_0^{(2)}\left(\frac{x}{2^{2l}}\right)}{\sqrt{2^2}} = \prod_{l=1}^{\infty} \frac{m_0\left(\frac{x}{2^{2l}}\right) m_0\left(\frac{x}{2^{2l-1}}\right)}{\sqrt{2^2}} = \widehat{\varphi}_{1,1}(x),$$

and similarly for $\widehat{\varphi}_{2,2}$. Hence, $\varphi_{1,1} = \varphi_{2,1} = \varphi_{2,2} =: \varphi = \frac{1}{3}\chi_{[0,3)}$, and $\varphi_0 = (\varphi, \varphi, \varphi)$.

To construct the wavelet we can pick

$$m_1(z) = \frac{1-z^3}{\sqrt{2}}, \quad (z \in \mathbb{T}).$$

Then the wavelet $\psi_0 = (\psi_1, \psi_2, \psi_3)$ is given by

$$U_0\psi_0 = \frac{1}{\sqrt{2}}(\varphi_0 - T_0^3\varphi_0)$$

so

$$\psi_1 = \psi_2 = \psi_3 =: \psi = \frac{1}{3}(\chi_{[0, \frac{3}{2})} - \chi_{[\frac{3}{2}, 1)}).$$

and

$$\{U_0^m T_0^n \psi_0 \mid m, n \in \mathbb{Z}\}$$

is an orthonormal basis for $L^2(\mathbb{R})^3$ which dilates the normalized tight frame of $L^2(\mathbb{R})$

$$\{U^m T^n \psi \mid m, n \in \mathbb{Z}\}.$$

Acknowledgements. The author wants to thank professor Șerban Strătilă for pointing out the connection between the GNS construction and Kolmogorov's theorem. This was the starting point and the key idea of this paper. Also many thanks to professor Palle Jorgensen for his suggestions and his constant support.

REFERENCES

- [BraJo97] O.Bratteli, P.E.T. Jorgensen, *Isometries, shifts, Cuntz algebras and multiresolution wavelet analysis of scale N* , Integral Equations Operator Theory, 28 (1997), 382-443
- [BraJo] O.Bratteli, P.E.T. Jorgensen, *Wavelets Through a Looking Glass*, book, to appear, Birkhauser.
- [BraRo] O.Bratteli, D.W.Robinson, *Operator Algebras and Quantum Statistical Mechanics*, vol. I, Springer-Verlag, 1987
- [C96] T. Constantinescu, *Schur Parameters, Factorization and Dilation Problems*, Operator Theory Advances and Applications, Vol. 82, Birkhauser Verlag, 1996
- [C97] T. Constantinescu, *Representations of Hermitian kernels in Krein spaces*, Publ. RIMS, Kyoto Univ, 33 (1997), 917-951
- [Dau92] I. Daubechies, *Ten Lectures on Wavelets*, CBMS-NSF Regional Conf. Ser. in Appl. Math., vol. 61, Society for Industrial and Applied Mathematics, Philadelphia, 1992.
- [Dut1] D. Dutkay, *Harmonic analysis of signed Ruelle transfer operators*, J. Math. Anal. Appl. 273 (2002), no. 2, 590-617
- [Dut2] D. Dutkay, *The wavelet Galerkin operator*, to appear in Journal of Operator Theory
- [EvKa] D.E. Evans, Y. Kawahigashi, *Quantum Symmetries on Operator Algebras*, Oxford Science Publications, 1998
- [EvLe] D.E. Evans, J.T. Lewis, *Dilations of irreversible evolutions in algebraic quantum theory*. Commun. Dubl. Inst. Adv. Studies, Ser. A., 24 (1977)
- [Fol] G. Folland, *A Course in Abstract Harmonic Analysis*, CRC Press, 1995
- [HL] D. Han, D. Larson, *Frames, bases and group representations*, Memoirs of the AMS, sept. 2000, vol.147, no. 697
- [HeWe] E. Hernandez, G. Weiss, *A First Course on Wavelets*, CRC Press, Inc. 1996
- [Jor98] P.E.T Jorgensen, *Ruelle operators : Functions which are harmonic with respect to a transfer operator*, Mem. Amer. Math. Soc.
- [Law91a] W.M. Lawton, *Necessary and sufficient conditions for constructing orthonormal wavelet bases*, J. Math. Phys. 32 (1991), 57-61
- [Sar67] D. Sarason, *generalized interpolation in H^∞* , Trans. Amer. Math. Soc., **127**(1967), 179-203
- [SzF68] B. Sz.-Nagy, C. Foias, *Dilatations des commutants d'operateurs*, C.R. Acad. Sci. Paris, Serie A, **266**(1968), 493-495
- [SzF70] B. Sz.-Nagy, C. Foias, *Harmonic Analysis of operators on Hilbert Space*, North Holland, Amsterdam-Budapest, 1970

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF IOWA, 14 MACLEAN HALL, IOWA CITY, IA 52242-1419, U.S.A.

E-mail address: ddutkay@math.uiowa.edu